

ORTHOGONAL POLYNOMIALS IN SEVERAL NON-COMMUTING VARIABLES. II

T. BANKS AND T. CONSTANTINESCU

ABSTRACT. In this paper we continue to investigate a certain class of Hankel-like positive definite kernels using their associated orthogonal polynomials. The main result of this paper is about the structure of this kind of kernels.

1. INTRODUCTION

Positive definite kernels are studied for their manifold applications. In this paper we consider a special type of kernels K defined on the free semigroup on N generators with the property that

$$K(\alpha\sigma, \tau) = K(\sigma, I(\alpha)\tau)$$

for any words α, σ, τ , where $I(\alpha)$ denotes the word obtained by writing α in the reverse order. These kernels appear in many situations, see for instance [6] and [7]. Our goal is to determine an explicit structure of the positive definite kernels satisfying the above invariance property. Since in the case $N = 1$ such kind of kernels are precisely the Hankel kernels, it is quite natural to consider associated orthogonal polynomials and to study their properties. Our main result establishes the connection between moments and Jacobi coefficients, as a multivariable extension of a classical result. We also describe the Jacobi coefficients of the free products of orthogonal polynomials.

2. ORTHOGONAL POLYNOMIALS

We introduce orthogonal polynomials on several hermitian variables and we discuss several general results. Especially, we emphasize the usefulness of a matrix notation that reduces very much the degree of complexity and makes clear the analogy with the classical, one-dimensional case. Let $\mathbb{F}_N^+$ be the unital free semigroup on N generators $1, \dots, N$ with lexicographic order $\prec$. In particular, $\mathbb{F}_1^+$ is the set $\mathbb{N}_0$ of nonnegative integers. The set of positive integers will be denoted by $\mathbb{N}$. The empty word is the identity element of $\mathbb{F}_N^+$ and the length of the word σ is denoted by $|\sigma|$. The length of the empty word is 0. There is a natural involution on $\mathbb{F}_N^+$ given by $I(i_1 \dots i_l) = i_l \dots i_1$ as well as a natural action of $\mathbb{F}_N^+$ on itself by juxtaposition, $(\sigma, \tau) \rightarrow \sigma\tau$, $\sigma, \tau \in \mathbb{F}_N^+$. Let $\mathcal{P}_N$ be the algebra of polynomials on N non-commuting indeterminates $X_1, \dots, X_N$ with complex coefficients. For any $\sigma = i_1 \dots i_l \in \mathbb{F}_N^+$, we define $X_\sigma = X_{i_1} \dots X_{i_l}$. Using this notation, each element $P \in \mathcal{P}_N$ can be uniquely written as

$$(2.1) \quad P = \sum_{\substack{\sigma \in \mathbb{F}_N^+ \\ 1}} c_\sigma X_\sigma,$$

with only finitely many coefficients c_σ different from zero. The length of the highest σ such that $c_\sigma \neq 0$ is the *degree* of P . We also have

$$P = \sum_{k \geq 0} P_k = \sum_{k \geq 0} \sum_{|\sigma|=k} c_\sigma X_\sigma,$$

where each P_k belongs to the vector space $\mathcal{L}_k^N$ of homogeneous polynomials of degree $k \geq 0$ in N variables $X_1, \dots, X_N$. The dimension of $\mathcal{L}_k^N$ is N^k . An involution $+$ on $\mathcal{P}_N$ can be introduced as follows: $X_k^+ = X_k$, $k = 1, \dots, N$; on monomials, $(X_\sigma)^+ = X_{I(\sigma)}$; in general, if P has the representation as in (2.1) then

$$(2.2) \quad P^+ = \sum_{\sigma \in \mathbb{F}_N^+} \bar{c}_\sigma X_\sigma^+,$$

and $\mathcal{P}_N$ is a unital, associative $*$ -algebra over $\mathbb{C}$. Let ϕ be a strictly positive functional on $\mathcal{P}_N$, that is, ϕ is a linear unital map on $\mathcal{P}_N$ and $\phi(P^+P) > 0$ for every $P \in \mathcal{P}_N - \{0\}$. The Gelfand-Naimark-Segal construction applied to ϕ gives a Hilbert space $\mathcal{H}_\phi$ such that $\{X_\sigma\}_{\sigma \in \mathbb{F}_N^+}$ is a linearly independent family in $\mathcal{H}_\phi$. The Gram-Schmidt procedure gives a family $\{\varphi_\alpha\}_{\alpha \in \mathbb{F}_N^+}$ of polynomials such that

$$(2.3) \quad \varphi_\alpha = \sum_{\beta \preceq \alpha} a_{\alpha,\beta} X_\beta, \quad a_{\alpha,\alpha} > 0;$$

$$(2.4) \quad \langle \varphi_\alpha, \varphi_\beta \rangle_\phi = \delta_{\alpha,\beta}, \quad \alpha, \beta \in \mathbb{F}_N^+,$$

where for $P_1, P_2 \in \mathcal{P}_N$,

$$\langle P_1, P_2 \rangle_\phi = \phi(P_2^+ P_1).$$

The elements φ_α , $\alpha \in \mathbb{F}_N^+$, will be called the orthonormal polynomials associated with ϕ . We notice that the use of the Gram-Schmidt process depends on the order chosen on $\mathbb{F}_N^+$. A different order would give a different family of orthogonal polynomials. Due to the natural grading on $\mathbb{F}_N^+$ it is possible to develop a base free approach to orthogonal polynomials. In the case of orthogonal polynomials on several commuting variables this is presented in [8]. However, in this paper we consider only the lexicographic order on $\mathbb{F}_N^+$.

The moments of ϕ are

$$s_\sigma = \phi(X_\sigma), \quad \sigma \in \mathbb{F}_N^+,$$

and we define the moment kernel of ϕ by the formula $K_\phi(\alpha, \beta) = s_{I(\alpha)\beta}$, $\alpha, \beta \in \mathbb{F}_N^+$. We notice that K_ϕ is a positive definite kernel on $\mathbb{F}_N^+$ and for $\alpha, \sigma, \tau \in \mathbb{F}_N^+$,

$$(2.5) \quad K_\phi(\alpha\sigma, \tau) = K_\phi(\sigma, I(\alpha)\tau).$$

This property can be viewed as a Hankel type condition. Conversely, it is easily seen that if a positive definite kernel K satisfies (2.5) then there exists a positive functional ϕ on $\mathcal{P}_N$ such that $K = K_\phi$.

2.1. Three term relations. Let ϕ be a unital strictly positive functional on $\mathcal{P}_N$ and let $\{\varphi_\alpha\}_{\alpha \in \mathbb{F}_N^+}$ be the orthonormal polynomials associated with ϕ . As in the commutative case (see [8]), it is very convenient to use a matrix notation, $\Phi_n = [\varphi_\sigma]_{|\sigma|=n}$ for $n \geq 0$ and $\Phi_{-1} = 0$. With this notation, the analogy with the classical

case $N = 1$ will be much more transparent. It turns out that the family $\{\Phi_n\}_{n \geq 0}$ satisfies a three-term recursive formula,

$$(2.6) \quad X_k \Phi_n = \Phi_{n+1} A_{n+1,k} + \Phi_n B_{n,k} + \Phi_{n-1} A_{n,k}^*,$$

for $k = 1, \dots, N$ and $n \geq 0$ (see [5] and [2]). Each matrix $B_{n,k}$, $n \geq 0$, $k = 1, \dots, N$, is a selfadjoint $N^n \times N^n$ matrix, while each $A_{n,k}$, $n > 0$, $k = 1, \dots, N$, is an $N^n \times N^{n-1}$ matrix such that

$$A_n = \begin{bmatrix} A_{n,1} & \dots & A_{n,N} \end{bmatrix}$$

is an upper triangular invertible matrix for every $n \geq 0$, with strictly positive elements on the diagonal. The fact that A_n is upper triangular comes from the lexicographic order that we use on $\mathbb{F}_N^+$. The invertibility of B_n is a consequence of the fact that ϕ is strictly positive and appears to be a basic translation of this information. The diagonal of A_n is strictly positive since we chose $a_{\alpha,\alpha} > 0$. A family $\mathcal{A} = \{A_{n,k}, B_{m,k} \mid n > 0, m \geq 0, k = 1, \dots, N\}$ of matrices satisfying all these properties will be called admissible. It turns out that there are no other restrictions on the matrices $A_{n,k}$, $B_{n,k}$ as shown by the following Favard type result mentioned in [5]. A similar result for the monic orthogonal polynomials, $p_\sigma = \frac{1}{a_{\sigma,\sigma}} \varphi_\sigma$ was recently mentioned in [1].

Theorem 2.1. *Let $\varphi_\sigma = \sum_{\tau \prec \sigma} a_{\sigma,\tau} X_\tau$, $\sigma \in \mathbb{F}_N^+$, be elements in $\mathcal{P}_N$ such that $\varphi_\emptyset = 1$ and $a_{\sigma,\sigma} > 0$. Assume that there exists an admissible family $\mathcal{A}$ of matrices such that for $k = 1, \dots, N$ and $n \geq 0$,*

$$X_k \Phi_n = \Phi_{n+1} A_{n+1,k} + \Phi_n B_{n,k} + \Phi_{n-1} A_{n,k}^*,$$

where $\Phi_n = [\varphi_\sigma]_{|\sigma|=n}$ for $n \geq 0$ and $\Phi_{-1} = 0$. Then there exists a unique strictly positive functional ϕ on $\mathcal{P}_N$ such that $\{\varphi_\sigma\}_{\sigma \in \mathbb{F}_N^+}$ is the family of orthonormal polynomials associated to ϕ .

There is also a family of Jacobi matrices associated with the three-term relation in the following way. For $P \in \mathcal{P}_N$ define

$$\Psi_\phi(P) \varphi_\sigma = P \varphi_\sigma.$$

Since the moment kernel has the Hankel type structure mentioned above in (2.5), it follows that each $\Psi_\phi(P)$ is a symmetric operator on the Hilbert space $\mathcal{H}_\phi$ with dense domain $\mathcal{P}_N$. Moreover, for $P, Q \in \mathcal{P}_N$,

$$\Psi_\phi(PQ) = \Psi_\phi(P) \Psi_\phi(Q),$$

and $\Psi_\phi(P) \mathcal{D} \subset \mathcal{D}$, hence Ψ_ϕ is an unbounded representation of $\mathcal{P}_N$. Also, $\phi(P) = \langle \Psi_\phi(P) 1, 1 \rangle_\phi$ for $P \in \mathcal{P}_N$. We distinguish the operators $\Psi_k = \Psi_\phi(X_k)$, $k = 1, \dots, N$, since $\Psi_\phi(\sum_{\sigma \in \mathbb{F}_N^+} c_\sigma X_\sigma) = \sum_{\sigma \in \mathbb{F}_N^+} c_\sigma \Psi_\sigma$. Let $\{e_1, \dots, e_N\}$ be the standard basis of $\mathbb{C}^N$ and define the unitary operator W from $l^2(\mathbb{F}_N^+)$ onto $\mathcal{H}_\phi$ such that $W(e_\sigma) = \varphi_\sigma$, $\sigma \in \mathbb{F}_N^+$. We see that $W^{-1} \mathcal{D}$ is the linear space $\mathcal{D}_0$ generated by e_σ , $\sigma \in \mathbb{F}_N^+$, so that we can define

$$J_k = W^{-1} \Psi_k W, \quad k = 1, \dots, N,$$

on $\mathcal{D}_0$. Each J_k is a symmetric operator on $\mathcal{D}_0$ and by (2.6), the matrix of (the closure of) J_k with respect to the orthonormal basis $\{e_\sigma\}_{\sigma \in \mathbb{F}_N^+}$ is

$$J_k = \begin{bmatrix} B_{0,k} & A_{1,k}^* & 0 & \dots \\ A_{1,k} & B_{1,k} & A_{2,k}^* & \\ 0 & A_{2,k} & B_{2,k} & \ddots \\ \vdots & & \ddots & \ddots \end{bmatrix}.$$

We call $(J_1, \dots, J_N)$ a Jacobi N -family on $\mathcal{D}_0$. It turns out that the usual admissibility conditions on $A_{n,k}$ and $B_{n,k}$ insure a joint model of a Jacobi family in the following sense.

Theorem 2.2. *Let $(J_1, \dots, J_N)$ be a Jacobi N -family and assume that the corresponding $\mathcal{A}$ is an admissible family of matrices. Then there exists a unique strictly positive functional ϕ on $\mathcal{P}_N$ with associated orthonormal polynomials $\{\varphi_\sigma\}_{\sigma \in \mathbb{F}_N^+}$ such that the map $W(e_\sigma) = \varphi_\sigma$, $\sigma \in \mathbb{F}_N^+$, extends to a unitary operator from $l^2(\mathbb{F}_N^+)$ onto $\mathcal{H}_\phi$ and $J_k = W^{-1}\Psi_k W$ for $k = 1, \dots, N$.*

For details about the proof of this result see [5].

2.2. Jacobi N -families and combinatorics of lattice paths. The matrices $A_{n,k}$ and $B_{n,k}$ contain the whole information about the orthonormal polynomials (or the moment kernel K_ϕ). Usually they are called the Jacobi coefficients of K_ϕ and can be calculated from the moments. For instance,

$$A_n = [a_{\alpha,\beta}]_{|\alpha|=|\beta|=n}^{-1} [a_{\alpha,\beta}]_{|\alpha|=|\beta|=n-1}^{\oplus N},$$

where $a_{\alpha,\beta}$ are the coefficients of the orthogonal polynomials and for a matrix A we use the notation

$$A^{\oplus l} = \underbrace{A \oplus \dots \oplus A}_{l \text{ times}}.$$

In their turn, the coefficients $a_{\alpha,\beta}$, $\beta \preceq \alpha$, can be calculated from the formula

$$a_{\alpha,\beta} = \frac{1}{\sqrt{D_{\alpha-1}D_\alpha}} \det [K(\alpha', \beta')]_{\alpha' \prec \alpha, \beta' \preceq \alpha, \beta' \neq \beta},$$

where

$$D_\alpha = \det [K(\alpha', \beta')]_{\alpha', \beta' \preceq \alpha}.$$

The formula for $B_{n,k}$ is somewhat more involved and we do not record it here. Instead we consider a different kind of relation between moments and Jacobi coefficients which appears to be more explicit. The case $N = 1$ is classical, see [9], [10]. The Jacobi N -family $(J_1, \dots, J_N)$ is a convenient tool to deal with this matter. Thus, for any $\sigma \in \mathbb{F}_N^+$ we have that

$$s_\sigma = \phi(X_\sigma) = \langle \Psi_\phi(X_\sigma)1, 1 \rangle_\phi = \langle \Psi_\sigma 1, 1 \rangle_\phi,$$

and by Theorem 2.2,

$$\langle \Psi_\sigma 1, 1 \rangle_\phi = \langle J_\sigma e_0, e_0 \rangle,$$

therefore we have

$$(2.7) \quad s_\sigma = \langle J_\sigma e_0, e_0 \rangle, \quad \sigma \in \mathbb{F}_N^+.$$

Now we introduce some special paths on $\mathbb{N}_0 \times \{1, \dots, N\} \times \mathbb{N}_0$. The allowed steps are the following: level steps $l_{n,m}^k$ from a point (n, k, m) to $(n+1, k, m)$, level steps $lp_{m,k}^n$ from a point (n, k, m) to (n, p, m) , for some $p \in \{1, \dots, N\} - \{k\}$, rise steps $r_{n,m}^k$ from a point (n, k, m) to $(n+1, k, m+1)$, and fall steps $f_{n,m}^k$ from a point (n, k, m) to $(n+1, k, m-1)$ (see Figure 1 for an example).




FIGURE 1. An example of a path for $N = 2$

We introduce a weight on steps by the formula

$$w(\text{step}) = \begin{cases} I & \text{if step} = lp_{m,k}^n \\ B_{m,k} & \text{if step} = l_{n,m}^k \\ A_{m+1,k} & \text{if step} = r_{n,m}^k \\ A_{m,k}^* & \text{if step} = f_{n,m}^k \end{cases}$$

where I denotes the identity matrix of appropriate size. If $\mathbf{p}$ is made of l steps, step 1, $\dots$, step l , then we define the weight of $\mathbf{p}$ by the formula

$$w(\mathbf{p}) = w(\text{step } l) \dots w(\text{step } 1).$$

Any word $\sigma \in \mathbb{F}_N^+ - \{\emptyset\}$ has a unique representation $\sigma = i_1^{k_1} \dots i_p^{k_p}$ with $1_1, \dots, i_p \in \{1, \dots, N\}$, $k_1, \dots, k_p > 0$, and $i_l \neq i_{l+1}$ for $l = 1, \dots, p-1$. We consider the set $\mathcal{M}_\sigma$ of all paths that start at $(0, i_p, 0)$ and end at $(|\sigma|, i_1, 0)$, with the property that the first k_p steps belong to $\mathbb{N}_0 \times \{i_p\} \times \mathbb{N}_0$, the next k_{p-1} steps belong to $\mathbb{N}_0 \times \{i_{p-1}\} \times \mathbb{N}_0$, and so on, until the last k_1 steps which belong to $\mathbb{N}_0 \times \{i_1\} \times \mathbb{N}_0$. These sets are related to the set of Motzkin paths. The Motzkin paths of length n are the paths in $\mathbb{N}_0^2$ made of level, fall, and rise steps, starting at $(0, 0)$ and ending at $(n, 0)$. Their set is denoted by $\mathcal{M}_n$ and the number of elements of $\mathcal{M}_n$ is given by the Motzkin number

$$M_n = \frac{1}{n} \sum_k \binom{n}{k} \binom{n-k}{k-1}.$$

It is easily seen that for any $\sigma \in \mathbb{F}_N^+ - \{\emptyset\}$ there is a bijection between $\mathcal{M}_\sigma$ and $\mathcal{M}_{|\sigma|}$. We can now describe a combinatorial structure of the moments.

Theorem 2.3. *Let ϕ be a strictly positive functional on $\mathcal{P}_N$ and let $\mathcal{A}$ be the admissible family of matrices associated with ϕ by (2.6). Then the moments of ϕ can be calculated by the formula*

$$(2.8) \quad s_\sigma = \sum_{\mathbf{p} \in \mathcal{M}_\sigma} w(\mathbf{p}), \quad \sigma \in \mathbb{F}_N^+ - \{\emptyset\}.$$

Proof. We consider the following points in $\mathbb{N}_0 \times \{1, \dots, N\} \times \mathbb{N}_0$:

$$P_{n,j} = (0, j, n), \quad n \geq 0, j = 1, \dots, N;$$

$$Q_{n,k,m} = (n, k, m), \quad m, n \geq 0, k = 1, \dots, N.$$

For $\sigma \in \mathbb{F}_N^+ - \{\emptyset\}$, $\sigma = i_1^{k_1} \dots i_p^{k_p}$, we claim that

$$J_\sigma = \begin{bmatrix} J_{0,0}^\sigma & J_{0,1}^\sigma & \cdots \\ J_{1,0}^\sigma & J_{1,1}^\sigma & \\ \vdots & & \ddots \end{bmatrix},$$

where the entry $J_{k,j}^\sigma$ gives the sum of (the weights of) the paths in $\mathbb{N}_0 \times \{1, \dots, N\} \times \mathbb{N}_0$ from P_{j,i_p} to $Q_{|\sigma|,i_1,k}$. The claim is clearly true for $|\sigma| = 1$ and then suppose it true for any word of length $\leq n$. Then consider a word σ of length $n+1$. Several cases can occur.

Case 1. $k = 0$. Let $\sigma = i_1^{k_1} \dots i_p^{k_p} = i_1 \tau$. First, assume $k_1 = 1$. Due to the fact that the level steps of type lp have weight I and by the induction hypothesis, we deduce that the sum of the paths from P_{j,i_p} to $Q_{|\sigma|,i_1,0}$ is

$$B_{0,i_1} J_{0,j}^\tau + A_{1,i_1}^* J_{1,j}^\tau,$$

which is precisely the $(0, j)$ entry of the product

$$J_{i_1} J_\tau = J_\sigma.$$

The case $k_1 > 1$ is similar, just by the induction hypothesis we deduce that the sum of paths from P_{j,i_p} to $Q_{|\sigma|,i_1,0}$ is again

$$B_{0,i_1} J_{0,j}^\tau + A_{1,i_1}^* J_{1,j}^\tau,$$

which is precisely the $(0, j)$ entry of the product

$$J_{i_1} J_\tau = J_\sigma.$$

Case 2. $j = 0$ is similar.

Case 3. $k, j \geq 1$, then the induction hypothesis implies that the sum of paths from P_{j,i_p} to $Q_{|\sigma|,i_1,k}$ is

$$A_{k,i_1} J_{k-1,j}^\tau + B_{k,i_1} J_{k,j}^\tau + A_{k+1,i_1}^* J_{k+1,j}^\tau,$$

which is precisely the (k, j) entry of J_σ . □

When all $B_{n,k}$ are zero, the level steps of type $l_{n,m}^k$ disappear, and our discussion is somewhat related to parts of [11].

Formula (2.8) can be also used to calculate the Jacobi coefficients from the moments in a relatively simple way. Let $\sigma \in \mathbb{F}_N^+ - \{\emptyset\}$. It is convenient to introduce the notation $i(n)$ in order to denote the n th letter of the word σ (from left to right). If $|\sigma| = 2n$, then there exists a unique path $\mathbf{p}_\sigma$ with corresponding weight

$$A_{1,i(i)}^* \dots A_{n,i(n)}^* A_{n,i(n)} \dots A_{1,i(i)}.$$

If $|\sigma| = 2n+1$, then there exists a unique path, still denoted $\mathbf{p}_\sigma$, with corresponding weight

$$A_{1,i(i)}^* \cdots A_{n,i(n)}^* B_{n+1,i(n+1)} A_{n,i(n)} \cdots A_{1,i(i)}.$$

In any case, let $\mathcal{M}_\sigma^* = \mathcal{M}_\sigma - \{\mathbf{p}_\sigma\}$. Also, we introduce the notation: $\tilde{A}_1 = A_1$, and for $n \geq 2$,

$$\tilde{A}_n = A_n A_{n-1}^{\oplus N} \cdots A_1^{\oplus N^{n-1}}.$$

Corollary 2.4. *The following formulae hold: for $n \geq 1$,*

$$A_n^* A_n = \left(\tilde{A}_{n-1}^* \right)^{-1} \left([K_\phi(\sigma, \tau)]_{|\sigma|=|\tau|=n} - \left[\sum_{\mathbf{p} \in \mathcal{M}_{I(\sigma)\tau}^*} w(\mathbf{p}) \right]_{|\sigma|=|\tau|=n} \right) \tilde{A}_{n-1}^{-1};$$

$$B_{0,k} = s_k, \quad k = 1, \dots, N,$$

and for $n \geq 1$, $k = 1, \dots, N$,

$$B_{n,k} = \left(\tilde{A}_n^* \right)^{-1} \left([K_\phi(k\sigma, \tau)]_{|\tau|=|\sigma|+2=n+1} - \left[\sum_{\mathbf{p} \in \mathcal{M}_{I(\sigma)\tau}^*} w(\mathbf{p}) \right]_{|\tau|=|\sigma|+2=n+1} \right) \tilde{A}_n^{-1}.$$

Due to the fact that A_n is an upper triangular matrix with strictly positive elements on the diagonal, the first relation of the previous result uniquely determine A_n by Cholesky factorization.

3. FREE PRODUCTS

The set $\mathcal{P}_N$ can be viewed as the free product of N copies of $\mathcal{P}_1$:

$$\mathcal{P}_N = \underbrace{\mathcal{P}_1 \star \dots \star \mathcal{P}_1}_{N \text{ times}} = \mathbb{C} \oplus \left(\oplus_{n \geq 1} \oplus_{i_1 \neq i_2, \dots, i_{n-1} \neq i_n} \mathcal{P}_{i_1}^0 \otimes \dots \otimes \mathcal{P}_{i_n}^0 \right),$$

where $\mathcal{P}_i^0$ is the set of polynomials in the variable X_i , $i = 1, \dots, N$, without constant term. This remark suggests that the simplest examples of families of orthogonal polynomials can be obtained by using free products. Some examples already appeared in [1]. Here we describe a general construction. This allows to introduce multivariable analogues of all classical orthogonal polynomials.

The simplest attempt to construct families of orthogonal polynomials on $\mathcal{P}_N$ would be to consider orthogonal polynomials associated with free products of positive functionals. Let ϕ_1 and ϕ_2 be two strictly positive functional on $\mathcal{P}_{N_1}$, respectively $\mathcal{P}_{N_2}$. Their free product $\phi = \phi_1 \star \phi_2$ on $\mathcal{P}_{N_1} \star \mathcal{P}_{N_2}$ is defined by $\phi(1) = 1$ and $\phi(P_{i_1} \dots P_{i_n}) = \phi_{i_1}(P_{i_1}) \dots \phi_{i_n}(P_{i_n})$ for $n \geq 1$, $i_1 \neq i_2, \dots, i_{n-1} \neq i_n$, $P_{i_k} \in \mathcal{P}_{N_{i_k}}^0$, and $i_k \in \{1, 2\}$ for $k = 1, \dots, n$. By results in [3], [4], ϕ is a positive functional. However, as it turns out, ϕ is not strictly positive. Thus, consider the case of two strictly positive functionals ϕ_1 and ϕ_2 on $\mathcal{P}_1$. Let K be the restriction of the moment kernel $K_{\phi_1 \star \phi_2}$ to the set $\{1, 12\}$. Its matrix is then

$$\begin{aligned} K &= \begin{bmatrix} \phi_1(X_1^2) & \phi_1(X_1^2)\phi_2(X_2) \\ \phi_2(X_2)\phi_1(X_1^2) & \phi_2(X_2)\phi_1(X_1^2)\phi_2(X_2) \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & \phi_2(X_2) \end{bmatrix} \begin{bmatrix} \phi_1(X_1^2) & \phi_1(X_1^2) \\ \phi_1(X_1^2) & \phi_1(X_1^2) \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & \phi_2(X_2) \end{bmatrix}. \end{aligned}$$

The matrix $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ has rank one, so K is never invertible.

We can consider another simple way related to free products in order to build orthogonal polynomials in several noncommutative variables. Thus, let $\{\varphi_{n,k}\}$, $n \geq 0$, $k \in \{1, \dots, N\}$, be N families of orthonormal polynomials on the real line, determined by the recursion formulae:

$$(3.1) \quad x\varphi_{n,k}(x) = a_{n+1,k}\varphi_{n+1,k}(x) + b_{n,k}\varphi_{n,k}(x) + a_{n,k}\varphi_{n-1,k}(x).$$

We introduce polynomials in N noncommutative variables as follows. Any word $\sigma \in \mathbb{F}_N^+ - \{\emptyset\}$ can be uniquely represented in the form $\sigma = i_1^{k_1} \dots i_p^{k_p}$, $i_1, \dots, i_p \in \{1, \dots, N\}$, $k_1, \dots, k_p > 0$, and $i_l \neq i_{l+1}$ for $l = 1, \dots, p-1$. Then define

$$(3.2) \quad \varphi_\sigma(X_1, \dots, X_N) = \varphi_{k_1, i_1}(X_{i_1}) \dots \varphi_{k_p, i_p}(X_{i_p}).$$

Theorem 3.1. *There exists an admissible family $\mathcal{A}$ of matrices such that for $k = 1, \dots, N$ and $n \geq 0$,*

$$(3.3) \quad X_k \Phi_n = \Phi_{n+1} A_{n+1,k} + \Phi_n B_{n,k} + \Phi_{n-1} A_{n,k}^*,$$

where $\Phi_n = [\varphi_\sigma]_{|\sigma|=n}$ for $n \geq 0$, $\Phi_{-1} = 0$, and φ_σ are given by (3.2).

Proof. First we prove the result for $N = 2$. From (3.1) we deduce

$$X_1 = \varphi_1 a_{1,1} + b_{0,1} = \begin{bmatrix} \varphi_1 & \varphi_2 \end{bmatrix} \begin{bmatrix} a_{1,1} \\ 0 \end{bmatrix} + b_{0,1}$$

so that

$$X_1 \Phi_0 = \Phi_1 A_{1,1} + \Phi_0 B_{0,1}$$

with

$$A_{1,1} = \begin{bmatrix} a_{1,1} \\ 0 \end{bmatrix} \quad \text{and} \quad B_{0,1} = \begin{bmatrix} b_{0,1} \end{bmatrix}.$$

Similarly,

$$X_2 \Phi_0 = \Phi_1 A_{1,2} + \Phi_0 B_{0,2}$$

with

$$A_{1,2} = \begin{bmatrix} a_{1,2} \\ 0 \end{bmatrix} \quad \text{and} \quad B_{0,2} = \begin{bmatrix} b_{0,2} \end{bmatrix}.$$

We see that

$$A_1 = \begin{bmatrix} A_{1,1} & A_{1,2} \end{bmatrix} = \begin{bmatrix} a_{1,1} & 0 \\ 0 & a_{1,2} \end{bmatrix}$$

is upper triangular (actually diagonal) and has the elements on the diagonal > 0 (since all $a_{n,k} > 0$). The case $n \geq 1$ can be dealt with in a similar manner. The first 2^{n-1} words of length n start with letter 1 and have the structure

$$1^{n-k} \tau$$

where τ is a word of length k starting with letter 2 (unless it is $\emptyset$). For $k = 0$ there is exactly one such word, 1^n , while for $0 < k \leq n-1$, there are 2^{k-1} such kind of words. Using (3.1) we have that

$$\begin{aligned} X_1 \varphi_{1^{n-k} \tau} &= X_1 \varphi_{1^{n-k}} \varphi_\tau = X_1 \varphi_{n-k,1} \varphi_\tau \\ &= \varphi_{n-k+1,1} \varphi_\tau a_{n-k+1,1} + \varphi_{n-k,1} \varphi_\tau b_{n-k,1} + \varphi_{n-k-1,1} \varphi_\tau a_{n-k,1} \\ &= \varphi_{1^{n-k+1} \tau} a_{n-k+1,1} + \varphi_{1^{n-k} \tau} b_{n-k,1} + \varphi_{1^{n-k-1} \tau} a_{n-k,1}. \end{aligned}$$

The last 2^{n-1} words of length n start with letter 2 and therefore we have for such a word σ that

$$\begin{aligned} X_1 \varphi_\sigma &= X_1 \varphi_{0,1} \varphi_\sigma \\ &= \varphi_{1,1} \varphi_\sigma a_{1,1} + \varphi_\sigma b_{0,1} \\ &= \varphi_{1\sigma} a_{1,1} + \varphi_\sigma b_{0,1}. \end{aligned}$$

Putting together the above two formulae we deduce that

$$X_1 \Phi_n = \Phi_{n+1} A_{n+1,1} + \Phi_n B_{n,1} + \Phi_{n-1} A_{n,1}^*,$$

where

$$A_{n,1} = \begin{bmatrix} a_{n,1} & & & & \\ & a_{n-1,1} & & & \\ & & a_{n-2,1}^{\oplus 2} & & \\ & & & \ddots & \\ & & & & a_{1,1}^{\oplus 2^{n-2}} \\ & & & & & 0_{2^{n-1} \times 2^{n-1}} \end{bmatrix}$$

and

$$B_{n,1} = \begin{bmatrix} b_{n,1} & & & & \\ & b_{n-1,1} & & & \\ & & b_{n-2,1}^{\oplus 2} & & \\ & & & \ddots & \\ & & & & b_{1,1}^{\oplus 2^{n-2}} \end{bmatrix};$$

the unspecified entries are all zero. Similarly, we deduce

$$X_2 \Phi_n = \Phi_{n+1} A_{n+1,2} + \Phi_n B_{n,2} + \Phi_{n-1} A_{n,2}^*,$$

for some suitable matrices $A_{n,2}$ and $B_{n,2}$. Actually, the same proof works for $N > 2$ and we record here the form of the matrices $A_{n,k}$, $B_{n,k}$ for an arbitrary N . Let

$$\mathcal{W}_k^n = \{\sigma \in \mathbb{F}_N^+ \mid |\sigma| = n \text{ and } \sigma = k\tau \text{ for some } \tau\}$$

and denote by π_k^n the bijection from the set $\{1, 2, \dots, N^{n-1}\}$ onto $\mathcal{W}_k^n$ defined simply by $\pi_k^n(l)$ = the l th word in $\mathcal{W}_k^n$, with respect to the lexicographic order. Also, if $\sigma \in \mathcal{W}_k^n$ then it has a unique representation $\sigma = k^p \tau$ with τ a word that does not start with the letter k . Define $n_k(\sigma) = p$. Now $B_{n,k}$ is an $N^n \times N^n$ matrix such that, for $l, m \in \{1, 2, \dots, N^n\}$,

$$(3.4) \quad (B_{n,k})_{m,l} = \begin{cases} b_{n_k(\pi_k^{n+1}(m)) - 1, k} & l = m \\ 0 & l \neq m \end{cases}$$

and $A_{n,k}$ is an $N^n \times N^{n-1}$ matrix such that for $l \in \{1, 2, \dots, N^{n-1}\}$ and $m \in \{1, 2, \dots, N^n\}$,

$$(3.5) \quad (A_{n,k})_{m,l} = \begin{cases} a_{n_k(\pi_k^n(m)) - 1, k} & l = m \\ 0 & l \neq m \end{cases}.$$

Therefore $A_n = \begin{bmatrix} A_{n,1} & \dots & A_{n,N} \end{bmatrix}$ is a diagonal matrix with strictly positive diagonal elements, so that $\mathcal{A} = \{A_{n,k}, B_{m,k} \mid n > 0, m \geq 0, k = 1, \dots, N\}$ is an admissible family of matrices. $\square$

REFERENCES

- [1] M. Anshelevich, Orthogonal polynomials with a resolvent-type generating function, arXiv, CO/0410482.
- [2] T. Banks, T. Constantinescu, and J. L. Johnson, Relations on noncommutative variables and associated orthogonal polynomials, arXiv, FA/0312389.
- [3] F. Boca, Completely positive maps on amalgamated product C^* -algebras, *Math. Scand.*, **72**(1993), 212-222.
- [4] M. Bożejko, Positive definite kernels, length functions on groups and noncommutative von Neumann inequality, *Studia Math.*, **95**(1989), 107-118.
- [5] T. Constantinescu, Orthogonal polynomials in several non-commuting variables. I, in *Spectral Theory and its Applications*, Theta, 2003, pp. 69-86.
- [6] T. Constantinescu and A. Gheondea, Representations of Hermitian kernels by means of Krein spaces II. Invariant kernels, *Commun. Math. Phys.*, **216**(2001), 409-430.
- [7] T. Constantinescu and A. Gheondea, On L. Schwartz's boundedness condition for kernels, *Positivity*, 2004, to appear.
- [8] C. H. Dunkl and Y. Xu, *Orthogonal Polynomials of Several Variables*, Cambridge Univ. Press, 2001.
- [9] P. Flajolet, Combinatorial aspects of continued fractions, *Discrete Math.*, **32**(1980), 125-161.
- [10] A. Lascoux, *Symmetric Functions and Combinatorial Operators on Polynomials*, CBMS, **99**, Amer. Math. Soc., Providence, Rhode Island, 2003.
- [11] A. Nica, R -transform of free joint distributions and non-crossing partitions, *J. Funct. Anal.*, **135**(1996), 271-296.
- [12] G. Szegő, *Orthogonal Polynomials*, Colloquium Publications, **23**, Amer. Math. Soc., Providence, Rhode Island, 1939.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS AT DALLAS, BOX 830688, RICHARDSON, TX 75083-0688, U. S. A.

E-mail address: banks@utdallas.edu

E-mail address: tiberiu@utdallas.edu